

Unsatisfiable CNF-formulas

Heidi Gebauer *

October 10, 2008

Abstract

A Boolean formula in a conjunctive normal form is called a (k, s) -formula if every clause contains exactly k variables and every variable occurs in at most s clauses. We show that there are unsatisfiable $(k, 4 \cdot \frac{2^k}{e})$ -CNF formulas.

1 A better bound for unsatisfiable formulas

Theorem 1.1 *For every sufficiently large k there is an unsatisfiable $(k, 4 \cdot \frac{2^k}{e})$ -CNF.*

Note that due to Kratochvíl, Savický and Tuza [2] every $(k, \frac{2^k}{e})$ -CNF is satisfiable. So our result shows that this bound is tight up to a factor $4e$.

Proof: We consider the class $\mathcal{C}$ of hypergraphs $\mathcal{G}$ whose vertices can be arranged in a binary tree $T_{\mathcal{G}}$ such that every hyperedge of $\mathcal{G}$ is a path of $T_{\mathcal{G}}$. For positive integers $k, s \geq 1$ we denote by a (k, s) -tree a k -uniform hypergraph $\mathcal{G} \in \mathcal{C}$ such that

- every full branch of $T_{\mathcal{G}}$ contains a hyperedge of $\mathcal{G}$ and
- every vertex of $T_{\mathcal{G}}$ belongs to at most s hyperedges of $\mathcal{G}$

When there is no danger of confusion we write $\mathcal{G}$ for $T_{\mathcal{G}}$. The following lemma is the core of our proof.

Lemma 1.2 *For every sufficiently large k there is a $(k, 2 \cdot \frac{2^k}{e})$ -tree $\mathcal{G}$.*

We first show that Lemma 1.2 implies Theorem 1.1. Suppose that there is a $(k, 2 \cdot \frac{2^k}{e})$ -tree $\mathcal{G}$ and let $\mathcal{G}'$ be a copy of $\mathcal{G}$. Let $\mathcal{H}$ be the hypergraph obtained by generating a new root v and attaching $\mathcal{G}$ as a left subtree and $\mathcal{G}'$ as a right subtree. Note that $\mathcal{H}$ is a $(k, 2 \cdot \frac{2^k}{e})$ -tree as well.

Let $(x_1, x'_1), (x_2, x'_2), \dots, (x_r, x'_r)$ denote the pairs of siblings of $\mathcal{H}$. We set $x'_i := \bar{x}_i$ for every i , $i = 1, \dots, r$ (i.e. each non-root vertex represents a literal $x \in \{x_1, \bar{x}_1, x_2, \bar{x}_2, \dots, x_r, \bar{x}_r\}$). Let $E(\mathcal{H})$

*Institute of Theoretical Computer Science, ETH Zurich, CH-8092 Switzerland. Email: gebauerh@inf.ethz.ch.

denote the set of hyperedges of $\mathcal{H}$. Then for every hyperedge $\{y_1, y_2, \dots, y_n\} \in E(\mathcal{H})$ we form the clause $C_{\{y_1, y_2, \dots, y_n\}} = (y_1 \vee y_2 \vee \dots \vee y_n)$ and set $\mathcal{F} := \bigwedge_{e \in E(\mathcal{H})} C_e$.

Note that every variable x_i of $\mathcal{F}$ occurs in at most $2 \cdot \Delta(F)$ clauses with $\Delta(F)$ denoting the maximum degree a variable in $\mathcal{F}$. Indeed, the number of occurrences of the variable x_i is bounded by the number of occurrences of the literal x_i plus the number of occurrences of the literal $\bar{x}_i$, which is at most $2\Delta F$. So $\mathcal{F}$ is a $(k, 2 \cdot \frac{2^k}{k})$ -CNF.

It remains to show that $\mathcal{F}$ is not satisfiable. Let α be an assignment to $\{x_1, \dots, x_r\}$.

Observation 1.3 *Note that there is (at least) one full branch b_{full} of $\mathcal{H}$ such that all literals along b_{full} are set to FALSE by α .*

By assumption b_{full} contains a hyperedge h . But α does not satisfy the clause C_h , implying that α does not satisfy $\mathcal{F}$. Since α was chosen arbitrarily, $\mathcal{F}$ is not satisfiable. $\square$

It remains to prove our key lemma.

Proof of Lemma 1.2: We need some notation first. The vertex set and the hyperedge set of a hypergraph $\mathcal{H}$ are denoted by $V(\mathcal{H})$ and $E(\mathcal{H})$, respectively. By a slight abuse of notation we consider $E(\mathcal{H})$ as a multiset, i.e. every hyperedge e can have a multiplicity greater than 1. By a *bottom hyperedge* of a tree $T_{\mathcal{H}}$ we denote a hyperedge covering a leaf of $T_{\mathcal{H}}$. Let $d = \frac{2^k}{k}$. For simplicity we assume that k is a power of 2, implying that d is power of 2 as well.

To construct the required hypergraph $\mathcal{G}$ we establish first a (not necessarily k -uniform) hypergraph $\mathcal{H}$ and then successively modify its hyperedges and $T_{\mathcal{H}}$. The following lemma is about the first step.

Lemma 1.4 *There is a hypergraph $\mathcal{H} \in \mathcal{C}$ with maximum degree $2d$ such that every full branch of $T_{\mathcal{H}}$ has 2^i bottom hyperedges of size $\log d + 1 - i$ for every i with $0 \leq i \leq \log d$.*

Proof of Lemma 1.4: Let T be a binary tree with $\log d + 1$ levels. In order to construct the desired hypergraph $\mathcal{H}$ we proceed for each vertex v of T as follows. For each leaf descendant w of v we let the path from v to w be a hyperedge of multiplicity $2^{l(v)}$ where $l(v)$ denotes the level of v . Figure 1 shows an illustration. The construction yields that each full branch of $T_{\mathcal{H}}$ has 2^i bottom hyperedges




Figure 1: An illustration of $\mathcal{H}$ for $d = 4$. The hyperedge $\{a, b, c\}$ has multiplicity 1, $\{b, c\}$ has multiplicity 2 and $\{c\}$ has multiplicity 4.

of size $\log d + 1 - i$ for every i with $0 \leq i \leq \log d$. So it remains to show that $d(v) \leq 2d$ for every vertex of $v \in V(T)$. Note that every vertex v has $2^{\log d - l(v)}$ leaf descendants in $T_{\mathcal{H}}$, implying that v is the start node of $2^{\log d - l(v)} \cdot 2^{l(v)} \leq d$ hyperedges. So the degree of the root is at most $d \leq 2d$. We then apply induction. Suppose that $d(u) \leq 2d$ for all nodes u with $l(u) \leq i - 1$ for some i with $1 \leq i \leq \log d$ and let v be a vertex on level i . By construction exactly half of the hyperedges containing the ancestor of v also contain v itself. Hence v occurs in at most $\frac{1}{2} \cdot 2d = d$ hyperedges as non-start node. Together with the fact that v is the start node of at most d hyperedges this implies that $d(v) \leq d + d \leq 2d$. $\square$

The next lemma deals with the second step of the construction of the required hypergraph $\mathcal{G}$.

Lemma 1.5 *There is a hypergraph $\mathcal{H}' \in \mathcal{C}$ with maximum degree $2d$ such that each full branch of $T_{\mathcal{H}'}$ has 2^i bottom hyperedges of size $\log d + 1 - i + \lfloor \log \log d \rfloor$ for some i with $0 \leq i \leq \log d$.*

Proof: Let $\mathcal{H} \in \mathcal{C}$ be a hypergraph with maximum degree $2d$ such that every leaf u of $T_{\mathcal{H}}$ is the end node of a set $S_i(u)$ of 2^i hyperedges of size $\log d + 1 - i$ for every i with $0 \leq i \leq \log d$. (Lemma 1.4 guarantees the existence of $\mathcal{H}$.) To each leaf u of $T_{\mathcal{H}}$ we then attach a binary tree T'_u of height $\lfloor \log \log d \rfloor$ in such a way that u is the root of T'_u . Let $v_0, \dots, v_{2^{\lfloor \log \log d \rfloor} - 1}$ denote the leaves of T'_u . For every i with $0 \leq i \leq 2^{\lfloor \log \log d \rfloor} - 1$ we then augment every hyperedge of $S_i(u)$ with the set of vertices different from u along the full branch of T'_u ending at v_i .

After repeating this procedure for every leaf u of $T_{\mathcal{H}}$ we get the desired hypergraph $\mathcal{H}'$. It remains to show that every vertex in $\mathcal{H}'$ has degree at most $2d$. To this end note first that during our construction the vertices of $\mathcal{H}$ did not change their degree. Secondly, let u be a leaf of $T_{\mathcal{H}}$. By assumption u has degree at most $2d$ and by construction $d(v) \leq d(u)$ for all vertices $v \in V(\mathcal{H}') \setminus V(\mathcal{H})$, which completes our proof. $\square$

Lemma 1.6 *There is a hypergraph $\mathcal{H}'' \in \mathcal{C}$ with maximum degree $2d$ such that every full branch of $T_{\mathcal{H}''}$ has one bottom hyperedge of size $\log d + 1 + \lfloor \log \log d \rfloor$.*

Note that due to our choice of d , Lemma 1.6 directly implies Lemma 1.2. $\square$

Proof of Lemma 1.6: By Lemma 1.5 there is a hypergraph $\mathcal{H}' \in \mathcal{C}$ with maximum degree $2d$ such that each full branch of $T_{\mathcal{H}'}$ has 2^i bottom hyperedges of size $\log d + 1 - i + \lfloor \log \log d \rfloor$ for some i with $0 \leq i \leq \log d$. For every leaf u of $T_{\mathcal{H}'}$ we proceed as follows. Let $e_1, \dots, e_{2^i}$ denote the bottom hyperedges of $\mathcal{H}'$ ending at u . We then attach a binary tree T'' of height i to u in such a way that u is the root of T'' . Let $p_1, \dots, p_{2^i}$ denote the full branches of T'' . We finally augment e_j with the vertices along p_j , for $j = 1 \dots 2^i$.

After repeating this procedure for every leaf u of $T_{\mathcal{H}'}$ we get the resulting graph $\mathcal{H}''$. By construction every full path of $T_{\mathcal{H}''}$ has one bottom hyperedge of size $\log d + 1 + \lfloor \log \log d \rfloor$. A similar argument as in the proof of Lemma 1.5 shows that the maximum degree of $\mathcal{H}''$ is at most $2d$. $\square$

References

- [1] N. Alon and J.H. Spencer, The Probabilistic Method *J. John Wiley & Sons* (2002).
- [2] J. Kratochvíl, P. Savický and Z. Tuza, One more occurrence of variables makes satisfiability jump from trivial to NP-complete *SIAM Journal of Computing* 22(1) **22(1)** (1993) 203210